

INDEX

- 5-hydroxytryptophan, and neuroendocrine responses in SAD patients, 46
- Alcoholism**, and light therapy, 78
- Alzheimer's disease, and light therapy, 78, 79
- American Sleep Disorders Association, 78
- Anorexia nervosa
See Eating disorders
- Antidepressants
See Medication treatment, of SAD
- Anxiety disorder
See Panic disorder
- Asia
prevalence of SAD, 37, 41
research studies on SAD, 26, 27
- Atenolol, as treatment for SAD, 43, 93
- Australia, genetic studies in incidence of SAD, 48
- B12**, as treatment for SAD, 93
- BDI
See Beck Depression Inventory (BDI)
- Beck Depression Inventory (BDI)
and diagnosis of SAD, 28, 32
and measurement of treatment effectiveness, 108
- Bipolar disorders
among SAD patients, 27, 33, 34-5, 36
and side effects of light therapy, 73
- Bulimia nervosa
See Eating disorders
- Bupropion, in treatment of SAD, 91
- Canadian Consensus Group** on seasonal affective disorder (SAD), 17, 18
Canadian Journal of Diagnosis, 19
- Canadian Psychiatric Association, 18
- Canadian Standards Association, 72
- Carbohydrates
craving for, in SAD patients, 23, 31, 71
and serotonin synthesis, 46
See also Eating disorders
- Catecholamine, depletion in SAD patients in remission with light therapy, 47
- Children
and light therapy, 76-7
with SAD, 44
- Chronic fatigue syndrome,
comorbidity with SAD, 106
- CIDI (Composite International Diagnostic Interview), 38
- Circadian rhythms
disorders, and light therapy, 64, 67, 68, 78-9
role in SAD, 43-5, 51
- Clarke Institute of Psychiatry (Toronto), 17
- Cocaine abuse, comorbidity with SAD, 106

- Cochrane Collaboration study, 65
- Cognitive-behavioural therapy (CBT), for SAD, 107-8
- Composite International Diagnostic Interview (CIDI), 38
- Concentration, inability for, 24
- D**-fenfluramine
 medication, as treatment for SAD, 93
 and serotonin transmission, 46
- Dawn simulation technique, 66-7
- Dementia, and light therapy, 78, 79
- Depression
 atypical, compared with SAD, 33-4
 nonseasonal, and light therapy, 77-8, 79
 nonseasonal, compared with SAD, 24, 25, 28
 severity, and choice of treatment, 96-8
 treatment resistant, 108-11
 vegetative symptoms, 23, 25
See also Light therapy; Medication treatment, of SAD; SAD (seasonal affective disorder)
- Depressive Experiences Questionnaire, 50
- Dexamethasone, and cortisol suppression in SAD patients, 48
- D,l-fenfluramine, and neuroendocrine responses in SAD patients, 46
- Dopamine, possible role in SAD, 47, 51
- DSM-III-R
 diagnostic criteria for SAD, 20, 21, 22, 23, 38-9
 personality disorders in SAD patients, and light therapy treatment outcome, 71
- DSM-IV, diagnostic criteria for SAD, 21, 23, 24, 33
- E**ating disorders
 appetite, increase in SAD patients, 23, 31, 33
 cognitive therapy for, 107
 comorbidity with SAD, 104, 105, 106, 107
 and light therapy, 78
 seasonal pattern, 35
See also Carbohydrates
- Elderly
 effect of light therapy on eyes, 75
 and insomnia, treatment with light therapy, 79
 and SAD, 40, 41
- Europe
 prevalence of SAD, 37, 41
 research studies on SAD, 26
- Eyes
 eyeblink, reduced in SAD patients, 47
 as receptors of light therapy, 68
 retina, role in SAD, 45, 51
 risk factors and light therapy, 73-5, 76
- F**atigue, in SAD patients, 48, 106
- Female-to-male ratio, of people with SAD, 39, 41
- Fenfluramine
 d-fenfluramine, medication, as treatment for SAD, 93
 d-fenfluramine, and serotonin transmission, 46
 d,l-fenfluramine, and neuroendocrine responses in SAD patients, 46
- Fluorescent light box
 availability, 72
 use in treatment of SAD, 64, 65-6, 67, 68, 69, 70, 76
- Fluoxetine, use in treatment of SAD, 89-90, 92, 96, 98, 101
- Freiburg Personality Inventory, 50
- G**enetic factors, in SAD, 47-8, 51
- Giessen Test, 50
- Global seasonality score (GSS), 28, 30, 31, 48
- GSS (global seasonality score), 28, 30, 31

Guilt, feelings of, 24

Hamilton Depression Rating Scale (HDRS)
and diagnosis of SAD, 28, 31-2
SIGH-SAD (Interview Guide), 32
and measurement of treatment effectiveness, 108

HDRS

See Hamilton Depression Rating Scale (HDRS)

Health and Welfare Canada, 17

HIGH-SAD (Hypomania Interview Guide for Seasonal Affective Disorder), 35

Hormonal changes, in SAD patients, 48-9

Hypericum, as treatment for SAD, 75, 93, 94

Hypersomnia, and SAD, 23, 24, 31, 33, 71

Hypomania

diagnostic instruments for, 34-5
Interview Guide for Seasonal Affective Disorder (HIGH-SAD), 35

side effect of light therapy, 73, 74

Hypothalamic-pituitary-adrenal (HPA) overactivity, in SAD patients, 48

ICD-IO, diagnostic criteria for SAD, 21

Iceland, lower SAD incidence compared with Americans, 48

Insomnia, and SAD, 24

Internet sites, on SAD, 19

Interpersonal psychotherapy (IPT), for SAD, 107-8

Inventory of Seasonal Variation (ISV), 31

ISV

See Inventory of Seasonal Variation (ISV)

Jet lag, and light therapy, 78, 79

Journal of Biological Rhythms, 78-9

L-tryptophan

medication, as treatment for SAD, 93, 94, 110, 111

and serotonin transmission, 46

Latitude, and prevalence estimate of SAD, 40, 41

LED light cap, 66

Levodopa, as treatment for SAD, 93

Light

cognitive sensitivity to, in SAD patients, 49-50

photoperiod decrease, and risk of SAD onset, 41, 51

Light therapy

definition, 64

devices, 64, 65-7, 72, 76

discontinuation of, 102, 104, 110

duration of exposure, 64, 65, 67, 68-9

intensity, 64, 65, 67, 68-9, 76, 81, 90, 93, 109

lights, incandescent versus halogen, 74, 76

melatonin rhythms, phase advance of, 43

parameters, 67-8

photoreceptors, 14-5, 45

placebo condition, appropriate, 80-1

positioning, and light source, 69, 76

response to, 33, 71, 109-10

reversal of, and neurotransmitter depletion, 47

risk factors, 73, 74, 75, 76

side effects, 73-5, 76, 98

during summer, 102-3, 104

timing, 44, 64, 67, 68, 69-70, 72, 76, 109

transdermal, 15, 68

treatment

decisions, 96-8

and medication, 75, 90, 99-101, 110, 111

for other disorders, 78-9, 107

outcome predictors, 71

practical tips for, 72-3

- trial length, 71, 76, 101
 - and tryptophan, effect reversal by deletion of, 46
 - wavelength, 67, 70, 73, 76
- Light visor, 66
- Longitudinal studies, of SAD patients
 - American (Leonhardt et al., Schwartz et al.), 26
 - British (Thompson et al.), 26
 - for identification of patients with bipolar disorders, 35
 - Japanese (Sakamoto et al., Sugishita et al.), 26, 27
 - Swiss (Graw et al.), 26
- M**-chlorophenylpiperazine, and neuroendocrine responses in SAD patients, 46
- Medical Research Council of Canada, 17
- Medication treatment, of SAD
 - combined with light therapy, 99-101
 - dosage, 91-2
 - side effects, 92
 - treatment decisions, 96-8
 - treatment trial length, 101
 - using
 - atenolol, 93
 - B12, 93
 - bupropion, 91
 - d-fenfluramine, 93
 - fluoxetine, 89-90, 92, 96, 98, 101
 - l-tryptophan, 93, 94, 110, 111
 - levodopa, 93
 - melatonin, 93, 94
 - moclobemide, 90-1
 - propranolol, 93
 - sertraline, 89, 90, 91-2, 101
 - tranylcypromine, 91
- Melatonin
 - medication, as treatment for SAD, 93, 94
 - role in SAD, 42-3, 51
 - suppression in light therapy, 64
- Metabolic changes, in SAD patients, 48-9
- Moclobemide, use in treatment of SAD, 90-1
- Mood reactivity, in SAD patients, 24, 33
- N**ational Comorbidity Survey (US), 38-9
- National Institute of Mental Health (NIMH), 13, 14, 26
- Neuroticism, in SAD patients, 50
- Neurotransmitters, role in SAD
 - dopamine, 47, 51
 - noradrenaline, 47, 51
 - serotonin, 14, 46
- Noradrenaline, possible role in SAD, 47, 51
- North America
 - higher incidence of SAD, compared with Icelanders, 48
 - prevalence of SAD, 37, 41
 - research studies on SAD, 26
- O**bsessive-compulsive disorder
 - comorbidity with SAD, 104, 105, 106
 - and light therapy, 78
- P**anic disorder
 - cognitive therapy for, 107
 - comorbidity with SAD, 104, 105, 106, 107
 - and light therapy, 78, 79
 - seasonal influence on, 35
- Personality disorders
 - comorbidity with SAD, 50, 51, 71, 105-6, 107
 - and light therapy, 78
- Pharmacological agents
 - See* Medication treatment, of SAD; names of specific drugs
 - "Phase shifting," treatment for SAD, 44
- Phototherapy
 - See* Light Therapy
- Placebo condition, for light treatment studies, 80-1

- Posttraumatic stress disorder, and
SAD, 35
- Premenopausal women, and SAD, 47
- Premenstrual syndrome
comorbidity with SAD, 35, 105, 107
and light therapy, 78, 79
- Prolactin, possible role in SAD, 47
- Propranolol, as treatment for SAD,
43, 93
- Psychological factors, in SAD pa-
tients, 49-50, 104-7
- Psychomotor impairment, in SAD
patients
agitation, 24
retardation/paralysis, 24, 33
- Psychotherapy, as treatment for SAD,
107-8, 111
- RDC** (*See* Research Diagnostic
Criteria), 34-5
- Remission
following treatment, 27
seasonal, 22-3
- Research Diagnostic Criteria (RDC),
34-5
- Retina, of SAD patients, 45
- Reverse vegetative symptoms, 23, 25,
33, 34
- Rosenthal, N.E.
diagnostic criteria for SAD, 21
light therapy as treatment for SAD,
14, 65
- S-SAD** (subsyndromal seasonal
affective disorder)
characteristics, 30
and light therapy, 77-8, 79
- SAD (seasonal affective disorder)
and circadian rhythms, 43-5, 51
compared with nonseasonal
depression, 25, 28, 33-5
consensus guidelines project, 17-9
demographic variables, 39, 40, 41
diagnosis stability, 25-8
diagnostic criteria
DSM-III-R, 20, 21, 22, 23
DSM-IV, 20, 23, 24, 33, 36
ICD-IO, 21
Rosenthal, 21
- diagnostic instruments
Beck Depression Inventory (BDI),
28, 32
Hamilton Rating Scale for Depres-
sion (HRSD), 28, 31-2
Inventory of Seasonal Variation
(ISV), 31
Seasonal Pattern Assessment
Questionnaire (SPAQ), 28-31, 36
- duration, 20-2
- epidemiology, 36-41
- etiology, 41-8
and eyes, role of, 45, 47, 51
genetic factors, 47-8, 51
historical references, 13
hormonal changes, 48-9
and latitude, effect of, 40, 41
and melatonin, role of, 42-3, 51
metabolic changes, 49
and neurotransmitters, role in, 46-
7, 51
onset, 20-2
pathophysiology, 48-51
personality factors, 50, 51, 71
photoperiod decrease, and risk of
onset, 41, 51
prevalence of, 36-41
psychological factors, 49-50
remission
following treatment, 27
seasonal, 22-3
research studies, 17-9, 25-8
screening for, 31, 37
and sex of patients, 39, 41
and thermoregulatory heat loss, 47
- treatment
antidepressant medication, 43,
89-94, 96-101, 110, 111
commencement of, 103-4
comorbid psychiatric disorders,
104-7
duration, 101-3, 104
high-density negative ions, 79, 80

- light therapy, 15, 33, 43, 64-79, 96-101
- limited response to, 108-11
- psychotherapy, 107-8
- relapse after discontinuation of, 101-2
- resistance to, 108-11
- sleep deprivation, 80
- walking exercise, 80, 107
- See also* Depression; Light therapy; Medication treatment, of SAD; S-SAD (subsyndromal seasonal affective disorder)
- Seasonal Pattern Assessment Questionnaire (SPAQ)
 - and genetic studies of SAD, 48
 - and determination of prevalence of SAD, 36-8
 - and diagnosis of SAD, 28-31
 - limitations of, 37-8
 - as screening instrument for SAD, 31, 37
- Serotonin, role in SAD, 14, 46
- Sertraline, use in treatment of SAD, 89, 90, 91-2, 101
- Sex, of patients with SAD, 39
- Shift work, and light therapy, 78, 79
- SIGH-SAD instrument
 - See* Hamilton Depression Rating Scale (HDRS)
- Sleep
 - See* Hypersomnia; Insomnia
- Society for Light Treatment and Biological Rhythms, 18, 78
 - formation of, 14
- Society for Research in Biological Rhythms, 18
- SPAQ
 - See* Seasonal Pattern Assessment Questionnaire (SPAQ)
- St. John's Wort, as treatment for SAD, 75, 93, 94
- Subsyndromal seasonal affective disorder
 - See* S-SAD (subsyndromal seasonal affective disorder)
- Suicidal ideation, 24
- Sumatriptan, and neuroendocrine responses in SAD patients, 46
- Summer remission, of depressive symptoms in SAD patients, 22-3
- Tanning salons**, 73
- Thermoregulatory heat loss, in SAD patients, 47
- Thyroid function, in SAD patients, 49, 51
- Tranlycypromine, use in treatment of SAD, 91
- Tryptophan
 - depletion in SAD patients in remission with light therapy, 47
 - and serotonin transmission, 46
- US National Comorbidity Survey**, 38-9
- US Underwriters Laboratory, 72
- Vegetative symptoms**, reverse, 23, 25, 33, 34
- Vitamin abnormalities, in SAD patients, 49
- Von Zerssen depression rating scale, 26
- Weight**
 - gain, 23, 24, 31, 33, 71
 - loss, 24
- Women
 - as majority of SAD patients, 39, 41
 - premenopausal, and SAD, 47
 - premenstrual syndrome, and SAD, 35